



Sedative-Hypnotics & the Treatment of Hypersomnia




Sedative-Hypnotics & the Treatment of Hypersomnia




- anxiolysis
- sedation-hypnosis
- anticonvulsant





Inhibition in the Brain



Inhibition in the Brain




- Today**
- GABA Receptors
 - Benzodiazepines
 - Barbituates
 - Amphetamines



5 Allosteric Modulation

definition: modulation achieved by binding of a drug to a site distinct from the site required for activation.

- Rudolph & Knoflach, 2011




types:

- positive (agonism)
 - benzodiazepines
- negative (inverse agonism)
 - βCCE

6

- antagonist (blocker)
 - Flumazenil

7



- Rudolph & Knoflach, 2011

Benzodiazepines

- there are many
 - Diazepam (*Valium*) among the first (launched 1963).
 - 4 benzodiazepines are among the 200 most commonly prescribed drugs in the U.S.
 - Alprazolam (*Xanax*)
 - Clonazepam (*Klonopin*)
 - Diazepam (*Valium*)
 - Lorazepam (*Ativan*)
- actions are dose-dependent:
 - most sedative hypnotics (e.g. barbituates)
 - death
 - anesthesia
 - hypnosis
 - sedation
 - anxiolysis

8

9

benzos by themselves do not:

- produce anesthesia
- cause fatalities

BUT
 they lower the lethal dose of other CNS depressants (e.g. alcohol)

from Patrice Guyenet, UVA Pharm Dept.

Benzodiazepines

- there are many
 - Diazepam (*Valium*) among the first (launched 1963).
 - 4 benzodiazepines are among the 200 most commonly prescribed drugs in the U.S.
 - Alprazolam (*Xanax*)
 - Clonazepam (*Klonopin*)
 - Diazepam (*Valium*)
 - Lorazepam (*Ativan*)
- actions are dose-dependent:
 - death
 - anesthesia
 - hypnosis
 - sedation
 - anxiolysis

Problems

- pharmacokinetics
- side effects

ideal hypnotic

ideal anxiolytic

from Patrice Guyenet, UVA Pharm Dept.

Benzodiazepines

- there are many
 - Diazepam (*Valium*) among the first (launched 1963).
 - 4 benzodiazepines are among the 200 most commonly prescribed drugs in the U.S.
 - Alprazolam (*Xanax*)
 - Clonazepam (*Klonopin*)
 - Diazepam (*Valium*)
 - Lorazepam (*Ativan*)
- actions are dose-dependent:
 - death
 - anesthesia
 - hypnosis
 - sedation
 - anxiolysis

Problems





- pharmacokinetics
- side effects

redistribution

metabolism

Benzodiazepines flurazepam

from Patrice Guyenet, UVA Pharm Dept.



Benzodiazepines: Last Couple of Things



- **Tolerance**
 - primarily observed with anticonvulsant actions
 - limited tolerance observed with sedative-hypnotic & anxiolytic effects
- **Dependence/Addiction**
 - physical dependence is usually mild
 - follows general rule of drug dependence:
 - higher dosage = more severe withdrawal
 - longer t1/2 = less severe withdrawal
 - estimated that 0.1-0.2% of adult population abuse or are dependent upon benzos (300,000-600,00 people in the U.S.)
 - GABA receptors live in the VTA (ventral tegmental area)
 - modulating GABA receptor activity in the VTA hypothesized to increase dopamine release
- **Benzodiazepine blocker**
 - Flumazenil (*Romazicon*)
 - benzodiazepine stupor
 - potential risk of seizures

Sedative-Hypnotics & the Treatment of Hypersomnia



Barbituates



- Directly bind to GABA binding site (at high doses)
 - activates channel and causes chloride conductance
- High doses are fatal
 - most sedative hypnotics (e.g. barbituates)
 - Benzodiazepines
- Once extensively used as sedative-hypnotics. Now largely replaced by the much safer benzos.
 - noteworthy exceptions:
 - Pentobarbital (insomnia, pre-op sedation, seizures)
 - Phenobarbital (seizures)
 - Thiopental (induction/maintenance of anesthesia)...short-lasting




Amphetamine



- Resembles catecholamines but more lipid soluble (can cross BBB)
- catecholamines: norepinephrine, dopamine, serotonin



Amphetamine




Ma huang
'looking for trouble'




- Resembles catecholamines but more lipid soluble (can cross BBB)
- catecholamines: norepinephrine, dopamine, serotonin
- indirectly-acting sympathomimetic amine
- amphetamine and related drugs stimulate release of:
 - dopamine → stimulates reward mechanisms, causes psychosis/addiction
 - norepinephrine → increased vigilance, anorexia
 - serotonin → increased vigilance, anorexia


CNS

sympathetic nerve terminals



- norepinephrine → hypertension, strokes, arrhythmias







- ### Amphetamine
- Powerful CNS stimulant
 - *d*-isomer 3-4 times more potent than *l*-isomer
 - *d*-amphetamine: Dextroamphetamine (*Dexedrine*, *Dextrostat*)
 - Lisdexamfetamine (*Vyvanse*): inactive, prodrug of *d*-amphetamine
 - Clinical uses:
 - Hypersomnia (Excessive Daytime Sleepiness [EDS])
 - narcolepsy (0.03-0.06% of the US population)
 - obstructive sleep apnea
 - shift-worker disorder (EDS affects >30% of night-shift workers)
 - Attention Deficit Hyperactivity Disorder
 - 23 ● Adverse/toxic effects
 - Usually result from overdosage
 - Acute toxic effects usually an extension of therapeutic effects.
 - restlessness, dizziness, tenseness, insomnia
 - Cardiovascular/GI side effects
 - Alternatives
 - Modafinil (*Provigil*): promotes wakefulness, reduces EDS in narcoleptics
 - 24 ● mechanism(s) not well-understood (but activates wake-promoting neurons)
 - little/no cardiovascular/cognitive side effects (main side effect = headaches)
 - may be used to reduce cocaine dependence



Sedative-Hypnotics & the Treatment of Hypersomnia

suggested reading

- **Basic & Clinical Pharmacology, 12th ed. (chapter 22)**
Bertram G. Katzung, Susan B. Masters, Anthony J. Trevor
- **Pharmacological Basis of Therapeutics, 12th ed. (Chapter 17)**
Goodman & Gilman

questions:
markbeen@virginia.edu

